arXiv:1501.00827v1 [math.FA] 5 Jan 2015

UNIVERSAL SERIES BY TRIGONOMETRIC SYSTEM IN WEIGHTED L^1_μ SPACES

S. A. EPISKOPOSIAN

Abstract. In this paper we consider the question of existence of trigonometric series universal in weighted $L^1_\mu[0, 2\pi]$ spaces with respect to rearrangements and in usual sense.

1. Introduction

Let X be a Banach space.

Definition 1.1. A series

$$\sum_{k=1}^{\infty} f_k, \quad f_k \in X \tag{1.1}$$

is said to be universal in X with respect to rearrangements, if for any $f \in X$ the members of (1.1) can be rearranged so that the obtained series $\sum_{k=1}^{\infty} f_{\sigma(k)}$ converges to f by norm of X.

Definition 1.2. The series (1.1) is said to be universal (in X) in the usual sense, if for any $f \in X$ there exists a growing sequence of natural numbers n_k such that the sequence of partial sums with numbers n_k of the series (1.1) converges to f by norm of X.

Definition 1.3. The series (1.1) is said to be universal (in X) concerning partial series, if for any $f \in X$ it is possible to choose a partial series $\sum_{k=1}^{\infty} f_{n_k}$ from (1.1), which converges to the f by norm of X.

1991 *Mathematics Subject Classification.* AMS Classification 2000 Primary 42A20 .

Key words and phrases. Universal series, trigonometric system, weighted spaces.

The author was supported in part by Grant- 01-000 from the Government of Armenia .

Note, that many papers are devoted (see [1]- [9]) to the question on existence of variouse types of universal series in the sense of convergence *almost everywhere and on a measure.*

The first usual universal in the sense of convergence almost everywhere trigonometric series were constructed by D.E.Menshov [1] and V.Ya.Kozlov [2]. The series of the form

$$\frac{1}{2}+\sum_{k=1}^{\infty} a_k coskx + b_k sinkx \tag{1.2}$$

was constructed just by them such that for any measurable on $[0,2\pi]$ function $f(x)$ there exists the growing sequence of natural numbers n_k such that the series (1.2) having the sequence of partial sums with numbers n_k converges to $f(x)$ almost everywhere on $[0,2\pi]$. (Note here, that in this result, when $f(x)\in L^1_{[0,2\pi]}$, it is impossible to replace convergence almost everywhere by convergence in the metric $L^1_{[0,2\pi]}$).

This result was distributed by A.A.Talalian on arbitrary orthonormal complete systems (see [3]). He also established (see [4]), that if $\{\phi_n(x)\}_{n=1}^{\infty}$ - the normalized basis of space $L^p_{[0,1]}, p>1$, then there exists a series of the form

$$\sum_{k=1}^{\infty} a_k\phi_k(x), \quad a_k\to 0. \tag{1.3}$$

which has property: for any measurable function $f(x)$ the members of series (1.3) can be rearranged so that the again received series converge on a measure on [0,1] to $f(x)$.

W. Orlicz [5] observed the fact that there exist functional series that are universal with respect to rearrangements in the sense of a.e. convergence in the class of a.e. finite measurable functions.

It is also useful to note that even Riemann proved that every convergent numerical series which is not absolutely convergent is universal with respect to rearrangements in the class of all real numbers.

Let $\mu(x)$ be a measurable on $[0,2\pi]$ function with $0<\mu(x)\le 1, x\in[0,2\pi]$ and let $L^1_{\mu}[0,2\pi]$ be a space of mesurable functions $f(x),\quad x\in[0,2\pi]$ with

$$\int_0^{2\pi}|f(x)|\mu(x)dx<\infty.$$

M.G.Grigorian constructed a series of the form (see [6]),

$$\sum_{k=-\infty}^{\infty} C_k e^{ikx} \quad with \quad \sum_{k=-\infty}^{\infty} |C_k|^q < \infty, \quad \forall q > 2$$

which is universal in $L^1_\mu[0,2\pi]$ concerning partial series for some weighted function $\mu(x), \quad 0 < \mu(x) \le 1, x \in [0,2\pi]$.

In [9] it is proved that for any given sequence of natural numbers $\{\lambda_m\}_{m=1}^{\infty}$ with $\lambda_m \nearrow^{\infty}$ there exists a series by trigonometric system of the form

$$\sum_{k=1}^{\infty} C_k e^{ikx}, \quad C_{-k} = \overline{C}_k, \tag{1.4}$$

with

$$\left|\sum_{k=1}^{m} C_k e^{ikx}\right| \le \lambda_m, \quad x \in [0,2\pi], \quad , \quad m = 1,2,...,$$

so that for each $\varepsilon > 0$ a weighted function $\mu(x)$,

$$0 < \mu(x) \le 1, |\{x \in [0,2\pi] : \mu(x) \ne 1\}| < \varepsilon$$

can be constructed, so that the series (1.4) is universal in the weighted space $L^1_\mu[0,2\pi]$ with respect simultaneously to rearrangements as well as to subseries.

In this paper we prove the following results.

Theorem 1.4. *There exists a series of the form*

$$\sum_{k=-\infty}^{\infty} C_k e^{ikx} \quad with \quad \sum_{k=-\infty}^{\infty} |C_k|^q < \infty, \quad \forall q > 2 \tag{1.5}$$

such that for any number $\varepsilon > 0$ *a weighted function* $\mu(x)$, $0 < \mu(x) \le 1$, *with*

$$|\{x \in [0,2\pi] : \mu(x) \ne 1\}| < \varepsilon \tag{1.6}$$

can be constructed, so that the series (1.5) *is universal in* $L^1_\mu[0,2\pi]$ *with respect to rearrangements .*

Theorem 1.5. *There exists a series of the form* (1.5) *such that for any number* $\epsilon > 0$ *a weighted function* $\mu(x)$ *with* (1.6) *can be constructed, so that the series* (1.5) *is universal in* $L^1_\mu[0,2\pi]$ *in the usual sense .*

2. BASIC LEMMA

Lemma 2.1. *For any given numbers* $0 < \varepsilon < \frac{1}{2}$, $N_0 > 2$ *and a step function*

$$f(x) = \sum_{s=1}^{q} \gamma_s \cdot \chi_{\Delta_s}(x), \tag{2.1}$$

where Δ_s *is an interval of the form* $\Delta_m^{(i)} = \left[\frac{i-1}{2^m}, \frac{i}{2^m}\right]$, $1 \le i \le 2^m$ *and*

$$|\gamma_s| \cdot \sqrt{|\Delta_s|} < \epsilon^3 \cdot \left(8 \cdot \int_0^{2\pi} f^2(x)dx\right)^{-1}, \quad s = 1, 2, ..., q. \tag{2.2}$$

there exists a measurable set $E \subset [0, 2\pi]$ *and a polynomial* $P(x)$ *of the form*

$$P(x) = \sum_{N_0 \le |k| < N} C_k e^{ikx}$$

which satisfy the conditions:

$$|E| > 2\pi - \varepsilon,$$

$$\int_E |P(x) - f(x)|dx < \varepsilon,$$

$$\sum_{N_0 \le |k| < N} |C_k|^{2+\varepsilon} < \varepsilon, \quad C_{-k} = \overline{C}_k$$

$$\max_{N_0 \le m < N} \left[\int_e \left|\sum_{N_0 \le |k| \le m} C_k e^{ikx}\right| dx\right] < \varepsilon + \int_e |f_(x)|dx,$$

for every measurable subset e *of* E.

Proof Let $0 < \epsilon < \frac{1}{2}$ be an arbitrary number.

Set

$$g(x) = 1, \quad if \quad x \in [0, 2\pi] \setminus \left[\frac{\varepsilon \cdot \pi}{2}, \frac{3\varepsilon \cdot \pi}{2}\right]; \tag{2.3}$$

$$g(x) = 1 - \frac{2}{\varepsilon}, \quad if \quad x \in \left[\frac{\varepsilon \cdot \pi}{2}, \frac{3\varepsilon \cdot \pi}{2}\right];$$

We choose natural numbers ν_1 and N_1 so large that the following inequalities be satisfied:

$$\frac{1}{2\pi}\left|\int_0^{2\pi} g_1(t)e^{-ikt}dt\right| < \frac{\varepsilon}{16 \cdot \sqrt{N_0}}, \quad |k| < N_0, \tag{2.4}$$

where

$$g_1(x) = \gamma_1 \cdot g(\nu_1 \cdot x) \cdot \chi_{\Delta_1}(x). \tag{2.5}$$

(By $\chi_E(x)$ we denote the characteristic function of the set E.) We put

$$E_1 = \{x \in \Delta_s : \quad g_s(x) = \gamma_s\}, \tag{2.6}$$

By (2.3), (2.5) and (2.6) we have

$$|E_1| > 2\pi \cdot (1-\epsilon) \cdot |\Delta_1|; \quad g_1(x) = 0, \quad x \notin \Delta_1, \tag{2.7}$$

$$\int_0^{2\pi} g_1^2(x)dx < \frac{2}{\epsilon} \cdot |\gamma_1|^2 \cdot |\Delta_1|. \tag{2.8}$$

Since the trigonometric system $\{e^{ikx}\}_{k=-\infty}^{\infty}$ is complete in $L^2[0,2\pi]$, we can choose a natural number $N_1 > N_0$ so large that

$$\int_0^{2\pi} \left| \sum_{0 \le |k| < N_1} C_k^{(1)} e^{ikx} - g_1(x) \right| dx \le \frac{\varepsilon}{8}, \tag{2.9}$$

where

$$C_k^{(1)} = \frac{1}{2\pi} \int_0^{2\pi} g_1(t) e^{-ikt} dt. \tag{2.10}$$

Hence by (2.4),(2.5) and (2.9) we obtain

(2.11)

$$\int_0^{2\pi} \left| \sum_{N_0 \le |k| < N_1} C_k^{(1)} e^{ikx} - g_1(x) \right| dx \le \frac{\varepsilon}{8} + \left[\sum_{0 \le |k| < N_0} |C_k^{(1)}|^2 \right]^{\frac{1}{2}} < \frac{\varepsilon}{4};$$

Now assume that the numbers $\nu_1 < \nu_2 < ...\nu_{s-1}$, $N_1 < N_2 < ... < N_{s-1}$, functions $g_1(x), g_2(x), ..., g_{s-1}(x)$ and the sets $E_1, E_2, ..., E_{s-1}$ are defined. We take sufficiently large natural numbers $\nu_s > \nu_{s-1}$ and $N_s > N_{s-1}$ to satisfy

$$\frac{1}{2\pi} \left| \int_0^{2\pi} g_s(t) e^{-ikt} dt \right| < \frac{\varepsilon}{16 \cdot \sqrt{N_{s-1}}}, \quad 1 \le s \le q, \quad |k| < N_{s-1}, \tag{2.12}$$

$$\int_0^{2\pi} \left| \sum_{0 \le |k| < N_s} C_k^{(s)} e^{ikx} - g_s(x) \right| dx \le \frac{\varepsilon}{4^{s+1}}, \tag{2.13}$$

where

$$g_s(x) = \gamma_s \cdot g(\nu_s \cdot x) \cdot \chi_{\Delta_s}(x), \qquad C_k^{(s)} = \frac{1}{2\pi} \int_0^{2\pi} g_s(t) e^{-ikt} dt. \tag{2.14}$$

Set

$$E_s = \{x \in \Delta_s : \quad g_s(x) = \gamma_s\}, \tag{2.15}$$

Using the above arguments (see (2.16)-(2.18)), we conclude that the function $g_s(x)$ and the set E_s satisfy the conditions:

$$|E_s| > 2\pi \cdot (1-\epsilon) \cdot |\Delta_s|; \quad g_s(x) = 0, \quad x \notin \Delta_s, \tag{2.16}$$

$$\int_0^{2\pi} g_s^2(x)dx < \frac{2}{\epsilon} \cdot |\gamma_s|^2 \cdot |\Delta_s|. \tag{2.17}$$

$$\int_0^{2\pi} \left| \sum_{N_{s-1} \le |k| < N_s} C_k^{(s)} e^{ikx} - g_1(x) \right| dx < \frac{\varepsilon}{2^{s+1}}. \tag{2.18}$$

Thus, by induction we can define natural numbers $\nu_1 < \nu_2 < ...\nu_q$, $N_1 < N_2 < ... < N_q$, functions $g_1(x), g_2(x), ..., g_q(x)$ and sets $E_1, E_2, ..., E_q$ such that conditions (2.14)- (2.16) are satisfied for all s, $1 \le s \le q$. We define a set E and a polynomial $P(x)$ as follows:

$$E = \bigcup_{s=1}^{q} E_s, \tag{2.19}$$

(2.20)

$$P(x) = \sum_{N_0 \le |k| < N} C_k e^{ikx} = \sum_{s=1}^{q} \left[\sum_{N_{s-1} \le |k| < N_s} C_k^{(s)} e^{ikx} \right], \quad C_{-k} = \overline{C}_k,$$

where

$$C_k = C_k^{(s)} \quad for \quad N_{s-1} \le |k| < N_s, \quad s = 1, 2, ..., q, \quad N = N_q - 1. \tag{2.21}$$

By Bessel's inequality and (2.3), (2.14) for all $s \in [1, q]$ we get

$$\left[\sum_{N_{s-1} \le |k| < N_s} |C_k^{(s)}|^2 \right]^{\frac{1}{2}} \le \left[\int_o^{2\pi} g_s^2(x)dx \right]^{\frac{1}{2}} \le \tag{2.22}$$

$$\le \frac{2}{\sqrt{\varepsilon}} \cdot |\gamma_s| \cdot \sqrt{|\Delta_s|}, \quad s = 1, 2, ..., q.$$

From (2.3), (2.12) and (2.13) it follows that

$$|E| > 2\pi - \varepsilon. \tag{2.23}$$

Taking relations (2.1), (2.3), (2.10), (2.12), (2.18) - (2.21) we obtain
(2.24)

$$\int_E |P(x)-f(x)|dx \le \sum_{s=1}^{q}\left[\int_E \left|\sum_{N_{s-1}\le|k|<N_s} C_k^{(s)}e^{ikx} - g_s(x)\right|dx\right] < \varepsilon$$

By (2.1), (2.2), (2.20) -(2.21)for any $k \in [N_0, N]$ we have

$$\sum_{N_0\le|k|<N} |C_k|^{2+\epsilon} \le \max_{N_0\le k\le N} |C_k|^{\epsilon} \cdot \sum_{k=N_0}^{N} |C_k|^2 \le \tag{2.25}$$

$$\le \max_{1\le s\le q}\left[\sqrt{\frac{8}{\epsilon}}\cdot|\gamma_s|\cdot\sqrt{|\Delta_s|}\right]\cdot\sum_{s=1}^{q}\left[\sum_{N_{s-1}\le|k|<N_s}|C_k^{(s)}|^2\right] \le$$

$$\le \max_{1\le s\le q}\left[\sqrt{\frac{8}{\epsilon}}\cdot|\gamma_s|\cdot\sqrt{|\Delta_s|}\right]\cdot\frac{8}{\epsilon}\cdot\sum_{s=1}^{q}|\gamma_s|^2\cdot|\Delta_s| \le$$

$$\le \max_{1\le s\le q}\left[\sqrt{\frac{8}{\epsilon}}\cdot|\gamma_s|\cdot\sqrt{|\Delta_s|}\right]\cdot\frac{8}{\epsilon}\cdot\left[\int_0^1 f^2(x)dx\right] < \epsilon;$$

That is, the statements 1) - 3) of Lemma are satisfied. Now we will check the fulfillment of statement 4) of Lemma. Let $N_0 \le m < N$, then for some s_0, $1 \le s_0 \le q$, $(N_{s_0} \le m < N_{s_0+1})$ we will have (see (2.20) and (2.21))
(2.26)

$$\sum_{N_0\le|k|\le m} C_k e^{ikx} = \sum_{s=1}^{s_0}\left[\sum_{N_{s-1}\le|k|<N_s} C_k^{(s)}e^{ikx}\right] + \sum_{N_{s_0-1}\le|k|\le m} C_k^{(s_0+1)}e^{ikx}.$$

Hence and from (2.1), (2.2), (2.3), (2.18), (2.19) and (2.22) for any measurable set $e \subset E$ we obtain

$$\int_e\left|\sum_{N_{s-1}\le|k|\le m} C_k e^{ikx}\right|dx \le$$

$$\le \sum_{s=1}^{s_0}\left[\int_e\left|\sum_{N_{s-1}\le|k|<N_s} C_k^{(s)}e^{ikx} - g_s(x)\right|dx\right] +$$

$$+\sum_{s=1}^{s_0}\int_e |g_s(x)|dx + \int_e\left|\sum_{N_{s_0-1}\le|k|\le m} C_k^{(s_0+1)}e^{ikx}\right|dx <$$

$$< \sum_{s=1}^{s_0}\frac{\varepsilon}{2^{s+1}} + \int_e |f(x)|dx + \frac{2}{\sqrt{\varepsilon}}\cdot|\gamma_{s_0+1}|\cdot\sqrt{|\Delta_{s_0+1}|} <$$

$$< \int_e |f(x)|dx + \varepsilon.$$

3. PROOF OF THEOREMS

Proof of Theorem 1.4 Let

$$f_1(x), f_2(x), ..., f_n(x), \quad x \in [0, 2\pi] \tag{3.1}$$

be a sequence of all step functions, values and constancy interval endpoints of which are rational numbers. Applying Lemma consecutively, we can find a sequence $\{E_s\}_{s=1}^{\infty}$ of sets and a sequence of polynomials

$$P_s(x) = \sum_{N_{s-1} \le |k| < N_s} C_k^{(s)} e^{ikx} \tag{3.2}$$

$$1 = N_0 < N_1 < ... < N_s < ...., \quad s = 1, 2, ...,$$

which satisfy the conditions:

$$|E_s| > 1 - 2^{-2(s+1)}, \quad E_s \subset [0, 2\pi], \tag{3.3}$$

$$\int_{E_s} |P_s(x) - f_s(x)|dx < 2^{-2(s+1)}, \tag{3.4}$$

$$\sum_{N_{s-1} \le |k| < N_s} \left| C_k^{(s)} \right|^{2+2^{-2s}} < 2^{-2s}, \quad C_{-k}^{(s)} = \overline{C}_k^{(s)} \tag{3.5}$$

$$\max_{N_{s-1} \le p < N_s} \left[\int_e \left| \sum_{N_{s-1} \le |k| \le p} C_k e^{ikx} \right| dx \right] < 2^{-2(s+1)} + \int_e |f_s(x)|dx, \tag{3.6}$$

for every measurable subset e of E_s.

Denote

$$\sum_{k=-\infty}^{\infty} C_k e^{ikx} = \sum_{s=1}^{\infty} \left[\sum_{N_{s-1} \le |k| < N_s} C_k^{(s)} e^{ikx} \right], \tag{3.7}$$

where $C_k = C_k^{(s)}$ for $N_{s-1} \le |k| < N_s$, $s = 1, 2, ...$

Let ε be an arbitrary positive number. Setting

$$\Omega_n = \bigcap_{s=n}^{\infty} E_s, \quad n = 1, 2, ...;$$

$$E = \Omega_{n_0} = \bigcap_{s=n_0}^{\infty} E_s, \quad n_0 = [\log_{1/2} \varepsilon] + 1; \tag{3.8}$$

$$B = \bigcup_{n=n_0}^{\infty} \Omega_n = \Omega_{n_0} \bigcup \left(\bigcup_{n=n_0+1}^{\infty} \Omega_n \setminus \Omega_{n-1} \right).$$

It is clear (see (3.3)) that $|B| = 2\pi$ and $|E| > 2\pi - \varepsilon$.

We define a function $\mu(x)$ in the following way:

$$\mu(x) = 1 \quad for \quad x \in E \cup ([0, 2\pi] \setminus B); \tag{3.9}$$

$$\mu(x) = \mu_n \quad for \quad x \in \Omega_n \setminus \Omega_{n-1}, \quad n \geq n_0 + 1,$$

where

$$\mu_n = \left[2^{4n} \cdot \prod_{s=1}^{n} h_s \right]^{-1}; \tag{3.10}$$

$$h_s = ||f_s(x)||_C + \max_{N_{s-1} \leq p < N_s} \| \sum_{N_{s-1} \leq |k| \leq p} C_k^{(s)} e^{ikx} \|_C + 1,$$

where

$$||g(x)||_C = \max_{x \in [0, 2\pi]} |g(x)|,$$

$g(x)$ is a continuous function on $[0, 2\pi]$.

From (3.5),(3.7)-(3.10) we obtain

(A) – $0 < \mu(x) \leq 1, \mu(x)$ is a measurable function and

$$|\{x \in [0, 2\pi] : \mu(x) \neq 1\}| < \varepsilon.$$

(B) – $\sum_{k=1}^{\infty} |C_k|^q < \infty, \quad \forall q > 2.$

Hence, obviously we have

$$\lim_{k \to \infty} C_k = 0. \tag{3.11}$$

It follows from (3.8)-(3.10) that for all $s \geq n_0$ and $p \in [N_{s-1}, N_s)$

$$\int_{[0, 2\pi] \setminus \Omega_s} \left| \sum_{N_{s-1} \leq |k| \leq p} C_k^{(s)} e^{ikx} \right| \mu(x) dx = \tag{3.12}$$

$$= \sum_{n=s+1}^{\infty} \left[\int_{\Omega_n \setminus \Omega_{n-1}} \left| \sum_{N_{s-1} \leq |k| \leq p} C_k^{(s)} e^{ikx} \right| \mu_n dx \right] \leq$$

$$\leq \sum_{n=s+1}^{\infty} 2^{-4n} \left[\int_0^{2\pi} \left| \sum_{N_{s-1}\leq|k|\leq p} C_k^{(s)} e^{ikx} \right| h_s^{-1} dx \right] < 2^{-4s}.$$

By (3.4), (3.8)-(3.10) for all $s \geq n_0$ we have

$$\int_0^{2\pi} |P_s(x) - f_s(x)| \, \mu(x)dx = \int_{\Omega_s} |P_s(x) - f_s(x)| \, \mu(x)dx+ \tag{3.13}$$

$$+\int_{[0,2\pi]\setminus\Omega_s} |P_s(x) - f_s(x)| \, \mu(x)dx = 2^{-2(s+1)}+$$

$$+\sum_{n=s+1}^{\infty} \left[\int_{\Omega_n\setminus\Omega_{n-1}} |P_s(x) - f_s(x)| \, \mu_n dx \right] \leq 2^{-2(s+1)}+$$

$$+\sum_{n=s+1}^{\infty} 2^{-4s} \left[\int_0^{2\pi} \left(|f_s(x)| + \left| \sum_{N_{s-1}\leq|k|<N_s} C_k^{(s)} e^{ikx} \right| \right) h_s^{-1} dx \right] <$$

$$< 2^{-2(s+1)} + 2^{-4s} < 2^{-2s}.$$

Taking relations (3.6), (3.8)- (3.10) and (3.12) into account we obtain that for all $p \in [N_{s-1}, N_s)$ and $s \geq n_0 + 1$

$$\int_0^{2\pi} \left| \sum_{N_{s-1}\leq|k|\leq p} C_k^{(s)} e^{ikx} \right| \mu(x)dx = \tag{3.14}$$

$$= \int_{\Omega_s} \left| \sum_{N_{s-1}\leq|k|\leq p} C_k^{(s)} e^{ikx} \right| \mu(x)dx+$$

$$+\int_{[0,2\pi]\setminus\Omega_s} \left| \sum_{N_{s-1}\leq|k|\leq p} C_k^{(s)} e^{ikx} \right| \mu(x)dx <$$

$$< \sum_{n=n_0+1}^{s} \left[\int_{\Omega_n\setminus\Omega_{n-1}} \left| \sum_{N_{s-1}\leq|k|\leq p} C_k^{(s)} e^{ikx} \right| dx \right] \cdot \mu_n + 2^{-4s} <$$

$$< \sum_{n=n_0+1}^{s} \left(2^{-2(s+1)} + \int_{\Omega_n\setminus\Omega_{n-1}} |f_s(x)| dx \right) \mu_n + 2^{-4s} =$$

$$= 2^{-2(s+1)} \cdot \sum_{n=n_0+1}^{s} \mu_n + \int_{\Omega_s} |f_s(x)| \mu(x)dx + 2^{-4s} <$$

$$< \int_0^{2\pi} |f_s(x)| \mu(x)dx + 2^{-4s}.$$

Let $f(x) \in L^1_\mu[0, 2\pi]$, i. e.$\int_0^{2\pi} |f(x)| \mu(x)dx < \infty$.

It is easy to see that we can choose a function $f_{\nu_1}(x)$ from the sequence (3.1) such that

$$\int_0^{2\pi} |f(x) - f_{\nu_1}(x)|\,\mu(x)dx < 2^{-2}, \quad \nu_1 > n_0 + 1. \tag{3.15}$$

Hence, we have

$$\int_0^{2\pi} |f_{\nu_1}(x)|\,\mu(x)dx < 2^{-2} + \int_0^{2\pi} |f(x)|\mu(x)dx. \tag{3.16}$$

From (2.1), (A), (3.13) and (3.15) we obtain with $m_1 = 1$

$$\int_0^{2\pi} \left|f(x) - \left[P_{\nu_1}(x) + C_{m_1}e^{im_1x}\right]\right| \mu(x)dx \leq \tag{3.17}$$

$$\leq \int_0^{2\pi} |f(x) - f_{\nu_1}(x)|\,\mu(x)dx +$$

$$+ \int_0^{2\pi} |f_{\nu_1}(x) - P_{\nu_1}(x)|\,\mu(x)dx +$$

$$+ \int_0^{2\pi} \left|C_{m_1}e^{im_1x}\right| \mu(x)dx < 2\cdot 2^{-2} + 2\pi\cdot |C_{m_1}|\,.$$

Assume that numbers $\nu_1 < \nu_2 < ... < \nu_{q-1}; m_1 < m_2 < ... < m_{q-1}$ are chosen in such a way that the following condition is satisfied:

$$\int_0^{2\pi} \left|f(x) - \sum_{s=1}^{j}\left[P_{\nu_s}(x) + C_{m_s}e^{im_sx}\right]\right| \mu(x)dx < \tag{3.18}$$

$$< 2\cdot 2^{-2j} + 2\pi\cdot \left|C_{m_j}\right|, \quad 1 \leq j \leq q-1.$$

We choose a function $f_{\nu_q}(x)$ from the sequence (3.1) such that

$$\int_0^{2\pi} \left|\left(f(x) - \sum_{s=1}^{q-1}\left[P_{\nu_s}(x) + C_{m_s}e^{im_sx}\right]\right) - f_{n_q}(x)\right| \mu(x)dx < 2^{-2q}, \tag{3.19}$$

where $\nu_q > \nu_{q-1}; \quad \nu_q > m_{q-1}$

This with (3.18) imply

$$\int_0^{2\pi} \left|f_{\nu_q}(x)\right| \mu(x)dx < 2^{-2q} + 2\cdot 2^{-2(q-1)} + 2\pi\cdot \left|C_{m_{q-1}}\right| = \tag{3.20}$$

$$= 9\cdot 2^{-2q} + 2\pi\cdot \left|C_{m_{q-1}}\right|.$$

By (3.13), (3.14) and (3.20) we obtain

$$\int_0^{2\pi} \left|f_{\nu_q}(x) - P_{\nu_q}(x)\right| \mu(x)dx < 2^{-2\nu_q}, \tag{3.21}$$

$$P_{\nu_q}(x)=\sum_{N_{\nu_q-1}\le|k|<N_{\nu_q}} C_k^{(\nu_q)}e^{ikx}.$$

(3.22)

$$\max_{N_{\nu_q-1}\le p<N\nu_q}\int_0^{2\pi}\left|\sum_{k=N_{\nu_q-1}}^{p} C_k^{(\nu_q)}e^{ikx}\right|\mu(x)dx<10\cdot 2^{-2q}+2\pi\cdot\left|C_{m_{q-1}}\right|.$$

Denote

$$m_q=\min\left\{n\in N: n\notin\left\{\left\{\{k\}_{k=N_{\nu_s-1}}^{N_{\nu_s}-1}\right\}_{s=1}^{q}\cup\{m_s\}_{s=1}^{q-1}\right\}\right\}. \tag{3.23}$$

From (2.1), (A), (3.19) and (3.21) we have

$$\int_0^{2\pi}\left|f(x)-\sum_{s=1}^{q}\left[P_{\nu_s}(x)+C_{m_s}e^{im_sx}\right]\right|\mu(x)dx\le \tag{3.24}$$

$$\le\int_0^{2\pi}\left|\left(f(x)-\sum_{s=1}^{q-1}\left[P_{\nu_s}(x)+C_{m_s}e^{im_sx}\right]\right)-f_{\nu_q}(x)\right|\mu(x)dx+$$

$$+\int_0^{2\pi}\left|f_{\nu_q}(x)-P_{\nu_q}(x)\right|\mu(x)dx+$$

$$+\int_0^{2\pi}\left|C_{m_q}e^{im_qx}\right|\mu(x)dx<2\cdot 2^{-2q}+2\pi\cdot\left|C_{m_q}\right|.$$

Thus, by induction we on q can choose from series (3.7) a sequence of members

$$C_{m_q}e^{im_qx},\quad q=1,2,...,$$

and a sequence of polynomials

$$P_{\nu_q}(x)=\sum_{N_{\nu_q-1}\le|k|<N_{\nu_q}} C_k^{(\nu_q)}e^{ikx},\quad N_{n_q-1}>N_{n_{q-1}},\quad q=1,2,.... \tag{3.25}$$

such that conditions (3.22) - (3.24) are satisfied for all $q\ge 1$.

Taking account the choice of $P_{\nu_q}(x)$ and $C_{m_q}e^{im_qx}$ (see (3.23) and (3.25)) we conclude that the series

$$\sum_{q=1}^{\infty}\left[\sum_{N_{\nu_q-1}\le|k|<N_{\nu_q}} C_k^{(\nu_q)}e^{ikx}+C_{m_q}e^{iqx}\right]$$

is obtained from the series (3.7) by rearrangement of members. Denote this series by $\sum C_{\sigma(k)}e^{i\sigma(k)x}$.

It follows from (3.11), (3.22) and (3.24) that the series $\sum C_{\sigma(k)}e^{i\sigma(k)x}$ converges to the function $f(x)$ in the metric $L^1_\mu[0,2\pi]$, i.e. the series (3.7) is universal with respect to rearrangements (see Definition 1.1).

The Theorem 1.4 is proved.

Proof of the Theorem 1.5

Applying Lemma consecutively, we can find a sequence $\{E_s\}_{s=1}^{\infty}$ of sets and a sequence of polynomials

$$P_s(x)=\sum_{N_{s-1}\le|k|<N_s} C_k^{(s)}e^{ikx},\quad C_{-k}^{(s)}=\overline{C}_k^{(s)} \tag{3.26}$$

$$1=N_0<N_1<...<N_s<....,\quad s=1,2,....,$$

which satisfy the conditions:

$$|E_s|>1-2^{-2(s+1)},\quad E_s\subset[0,2\pi], \tag{3.27}$$

$$\sum_{N_{s-1}\le|k|<N_s}\left|C_k^{(s)}\right|^{2+2^{-2s}}<2^{-2s}, \tag{3.28}$$

$$\int_{E_n}\left|f_n(x)-\sum_{s=1}^{n}P_s(x)\right|dx<2^{-n},\quad n=1,2,..., \tag{3.29}$$

where $\{f_n(x)\}_{n=1}^{\infty}$, $x\in[0,2\pi]$ be a sequence of all step functions, values and constancy interval endpoints of which are rational numbers.

Denote

$$\sum_{k=-\infty}^{\infty}C_ke^{ikx}=\sum_{s=1}^{\infty}\left[\sum_{N_{s-1}\le|k|<N_s}C_k^{(s)}e^{ikx}\right], \tag{3.30}$$

where $C_k=C_k^{(s)}$ for $N_{s-1}\le|k|<N_s$, $s=1,2,....$

It is clear (see (3.28)) that

$$\sum_{k=1}^{\infty}|C_k|^q<\infty,\quad \forall q>2.$$

Repeating reasoning of Theorem 1 a weighted function $\mu(x)$, $0<\mu(x)\le 1$ can constructed so that the following condition is satisfied:

$$\int_0^{2\pi}\left|f_n(x)-\sum_{s=1}^{n}P_s(x)\right|\cdot\mu(x)dx<2^{-2n},\quad n=1,2,... \tag{3.31}$$

For any function $f(x) \in L^1_\mu[0,1]$ we can choose a subsystem $\{f_{n_\nu}(x)\}_{\nu=1}^\infty$ from the sequence (3.1) such that

$$\int_0^{2\pi} |f(x) - f_{n_\nu}(x)|\, \mu(x)dx < 2^{-2\nu}. \tag{3.32}$$

From (3.30)-(3.32) we conclud

$$\int_0^{2\pi} \left| f(x) - \sum_{|k|\le M_\nu} C_k e^{ikx} \right| \mu(x)dx =$$

$$\int_0^{2\pi} \left| f(x) - \sum_{s=1}^{n_\nu} \left[\sum_{N_{s-1}\le|k|<N_s} C_k^{(s)} e^{ikx} \right] \right| \mu(x)dx \le$$

$$\le \int_0^{2\pi} |f(x) - f_{\nu_k}(x)| \cdot \mu(x)dx +$$

$$+ \int_0^{2\pi} \left| f_{\nu_k}(x) - \sum_{s=1}^{\nu_k} P_s(x) \right| \cdot \mu(x)dx < 2^{-2k} + 2^{-2\nu_k}$$

where $M_\nu = N_{n_\nu} - 1$.

Thus, the series (3.30) is universal in $L^1_\mu[0,1]$ in the sense of usual (see Definition 1.2).

The Theorem 1.5 is proved.

The author thanks Professor M.G.Grigorian for his attention to this paper.

References

1. D.E.Menshov, *On the partial summs of trigonometric series.*Mat. Sb. 20(1947), 197-238
2. V.Ya.Kozlov, *On the complete systems of orthogonal functions*, Mat. Sb. 26(1950), 351-364.
3. A.A.Talalian , *On the convergence almost everywhere the sumbsequence of partial summs of general orthogonal series*, Izv. Ak. Nauk Arm. SSR ser. Math. 10(1957), 17-34.
4. —— , *On the universal series with respect to rearrangements*, Izv. AN. SSSR ser. Math. 24(1960), 567-604.
5. W.Orlicz, *Uber die unabhangig von der Anordnung fast uberall kniwergenten Reihen*, Bull. de l'Academie Polonaise des Sciences, 81 (1927), p. 117-125;
6. M. G. Grigorian , *On the representation of functions by orthogonal series in weighted L^p spaces*, Studia. Math. 134(3)1999, 211-237.
7. P.L.Ul'janov , *On the unconditional convergence and the summmability*, Izv. AN. SSSR ser. Math. 22(1958), 811-840;
8. F.G.Arutunian, *To representation of functions by multiplaer series*D.A.N. Arm. SSR 64(1976), 72-76;
9. S.A.Episkoposian, M.G.Grigorian, *Representation of functions in weighted spaces $L^1_\mu[0, 2\pi]$ by trigonometric and Walsh series* , Analysis Mathematica, 27 (2001), 261-277;

DEPARTMENT OF PHYSICS, CHAIR OF HIGHER MATHEMATICS, YEREVAN STATE UNIVERSITY, YEREVAN, AL. MANOOGIAN ST.1, 375049, ARMENIA